CAS 79-94-7

Substance name

2,2',6,6'-tetrabromo-4,4'-isopropylidenediphenol (TBBPA)

Toxicity

An oral study in pregnant rats with TBBPA in its formulated product, Saytex 111, reported reduced fetal weight, increased malformations, and fetal death (ICI Americas 1985 study cited in NIEHS¹). Multiple subsequent studies on the technical compound did not show consistent reproductive or developmental toxicity. Kidney toxicity following oral dosing was reported in newborn rats.

TBBPA has been shown to compete with thyroid hormone (T4) in binding to transthyretin serum binding protein *in vitro*.^{1,2} It also appears to have potential to act as a thyroid hormone antagonist.^{4,5} TBBPA binds to the estrogen receptor but does not appear to be a receptor agonist or to have significant estrogenic potential.^{2,6} It is not currently listed as an endocrine disruptor by the European Union.

Exposure

This substance is listed as a Persistent, Bioaccumulative and Toxic (PBT) chemical under Washington State's PBT rule (WAC 173-333-320).⁷ TBBPA has been detected in breast milk in several small studies of the general population in Europe.^{2,8}

TBBPA is a high production volume (HPV) chemical that is used as both a reactive and additive flame retardant in plastics, adhesives, paper, and textiles.^{1,2,} It may constitute up to 22% of ABS polymer resins.² It is used primarily in electrical and electronic equipment. TBBPA is also used as a plasticizer, a component in adhesives and coatings, and a chemical intermediate for the synthesis of other flame retardants (e.g., TBBPA allyl ether).²

References

- National Institute of Environmental Sciences, Research Triangle Park. Tetrabromobisphenol A: Review of Toxicological Literature. June 2002. http://ntp.niehs.nih.gov/ntp/htdocs/Chem Background/ExSumPdf/tetrabromobisphenola.pdf.
- European Commission, European Chemicals Bureau. European Union Risk Assessment Report for 2,2',6,6'-tetrabromo-4,4'-isopropylidenediphenol (tetrabromobisphenol-A or TBBP-A) Part II – human health. 2006 Final Report. http://ecb.jrc.ec.europa.eu/DOCUMENTS/Existing-Chemicals/RISK_ASSESSMENT/REPORT/tbbpaHHreport402.pdf.
- 3. Fukuda, N et al. (2004) Unexpected nephrotoxicity induced by tetrabromobisphenol A in newborn rats. *Toxicol. Lett.* 150 (2): 145-155.
- Kitamura, S et al. (2005) Anti-thyroid hormone activity of tetrabromobisphenol A, a flame retardant, and related compounds: Afinity to the mammalian thyroid hormone receptor, and effect on tadpole metamorphosis. *Life Sci.* 76(14):1589-601.
- 5. Sun, H et al. (2009) Anti-thyroid hormone activity of bisphenol A, tetrabromobisphenol A and tetrachlorobisphenol A in an improved reporter gene assay. *Toxicol. In Vitro* 23(5): 950-4.
- 6. Dorosh, A et al. (2011) Assessing oestrogenic effects of brominated flame retardants hexabromocyclododecane and tetrabromobisphenol A on MCF-7 cells. *Folia. Biol.* 57 (1):35-9.

Rationale for Reporting List of Chemicals of High Concern to Children

Prepared by the Washington State Department of Health for the Children's Safe Product Act – 4/18/2011

- 7. WA Department of Ecology. Summary of Technical Background Information for the Proposed PBT List (Revised Draft) October 2005.
- Abdallah, MA and S. Harrad (2011) Tetrabromobisphenol-A, hexabromocyclododecane and its degradation products in UK human milk: relationship to external exposure. *Environ. Int. 37*(2): 443-8.